

MAVRIK

AQUAFLOW®

FOR COMMERCIAL USE ONLY ON INDOOR PLANTS IN GREENHOUSES AND INTERIOR PLANTSCAPES, OUTDOOR ORNAMENTAL PLANTINGS AND PLANTSCAPES, INCLUDING TREES, SHRUBS, AND FLOWER AND FOLIAGE CUTTINGS

SPECIMEN LABEL

ACTIVE INGREDIENT:

Tau-fluvalinate (CAS #102851-06-9) . . . 22.3%

OTHER INGREDIENTS: 77.7%

Total . . . 100.0%

MAVRIK AQUAFLOW® is a flowable formulation containing 2 pounds tau-fluvalinate per gallon (240 g/L).

EPA Reg. No. 2724-478

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Additional Precautionary Statements and First Aid

**SHAKE WELL BEFORE USING
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. **Avoid Breathing Spray Mist.** Certain persons may be sensitive to MAVRIK AQUAFLOW®'s fine spray particles. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

Sensitive individuals may temporarily experience an itching, burning, or tingling sensation, with or without a rash, following exposure. These symptoms will usually subside without requiring medical treatment. Avoid hand or sleeve-to-face contact. Prior to exposure or after washing, an application of corn oil to exposed skin may reduce these symptoms. Remove contaminated clothing and wash before reuse.

FIRST AID

IF INHALED

Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES

Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN

Wash with plenty of soap and water. Apply corn oil to affected area to relieve symptoms. Get medical attention if irritation or symptoms develop.

IF SWALLOWED

Call a physician or poison control center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with a finger. Do not induce vomiting or give anything by mouth to an unconscious person.

For 24 hour medical or chemical emergency assistance

Call Wellmark International (1-800-248-7763)

PERSONAL PROTECTIVE EQUIPMENT (PPE)

WPS USES:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) - in general, agricultural-plant uses are covered - must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, butyl rubber (≥14 mils), nitrile rubber (≥14 mils), neoprene rubber (≥14 mils), polyvinyl chloride (PVC) (≥14 mils), or viton (≥14 mils)
- Shoes plus socks
- For exposures outdoors, dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any R, P, or HE filter

USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, or viton
- Shoes plus socks
- Protective eyewear

INSTRUCTIONS FOR USE

Partially fill spray tank with water, then add recommended amount of ENSTAR® II, agitate, and complete filling.

Diluted spray solution should be used within 24 hours; always agitate before using.

RECOMMENDED GREENHOUSE APPLICATIONS RATE OF APPLICATION

FOR WHITEFLIES, APHIDS, SOFT-BODIED AND ARMORED SCALES

Preventative Program: Before insect populations reach damaging levels, mix ENSTAR II at the rate of 5 ounces per 100 gallons of water (1/3 tsp/1 gal) and apply to runoff every 2 weeks.

Curative Program: For an existing population of adult insects, make 2 applications 7 days apart of ENSTAR II to runoff at a rate of 10 ounces per 100 gallons of water (1 tsp/gal). Two weeks later, begin the Preventative Program. 100 gallons of mixture should be sprayed over 15,000 to 20,000 square feet.

FOR FUNGUS GNAT CONTROL

For the control of larvae and adult fungus gnats, apply ENSTAR II at 5 oz per 100 gallons of water (1.5 tsp/5 gal/1,000 sq ft) for moderate infestations to 10 oz (3 tsp/5 gal/1,000 sq ft) for heavy infestations. Complete coverage of soil is essential for fungus gnat control. Water as a drench after spraying to insure ENSTAR II contact with larvae and eggs. ENSTAR II may be applied for fungus gnat control before or after emergence of seedlings or on established production. Make 2 applications 7 to 10 days apart. Monitor fungus gnat population and make additional applications, if necessary.

FOR ROOT (GROUND) MEALYBUG

When infestations of root mealybugs are observed, the entire pot containing the infested plant should be drenched by submersion in a solution of ENSTAR II at a rate of one-half (1/2) ounce per 5 gallons of water [3 tsp/5 gal]. Submersion should continue until no more air bubbles are seen escaping from the fluid surface. This insures that the root system is saturated and the container is thoroughly treated. Repeat as necessary.

SQUARE FOOTAGE

When using 5 oz of ENSTAR II for Preventative Program or 10 oz for a Curative Program, refer to table below for square footage calculations:

Water to be Used*	100 gal	50 gal	25 gal	12.5 gal	5 gal
Square Ft	20,000	10,000	5,000	2,500	1,000
Preventative	5 oz	2 1/2 oz	6 1/2 tsp	3 1/4 tsp	1 1/2 tsp
Curative	10 oz	5 oz	12 tsp	6 tsp	3 tsp

* When using Electrostatic Spraying System (ESS®) or Puls-fog® equipment, apply the recommended amount of ENSTAR II over the square footage with the manufacturer's requirement of water. Follow manufacturer's calibration procedures.

METHODS OF APPLICATION

Do not apply ENSTAR II through any type of irrigation system.

Complete coverage of foliage is essential for control of whiteflies, aphids, mealybugs, fungus gnats, and scales. For conventional spray equipment, mix the recommended amount of ENSTAR II and water and spray both top and bottom surfaces of leaves until runoff. For other applications, follow the equipment manufacturer's specifications.

ENSTAR II is compatible with other commonly used insecticides and miticides, such as MAVRIK AQUAFLOW®. Before mixing in spray tank; however, it is advisable to test physical compatibility by mixing all components in a small container in proportionate quantities. Read and follow all precautionary statements, mixing instructions, and label directions on tank mix products.

EFFECTS ON GREENHOUSE PLANTS

Slight to moderate injury has occasionally been noted on some blooms under certain conditions. For insect control, it is suggested that applications be made in the pre-bloom stage. Before making large-scale application to any crop, a grower should test spray small numbers of plants to determine plant safety under his own cultural conditions.

Applications of ENSTAR II have been made to greenhouse plants (aluminum plant, azalea, Boston fern, bromeliad, chrysanthemum, devil's ivy, dianthus, dracaena compacta, emerald gem nephthylis, English ivy, epipremnum aureum, exotica perfection dumbcane, false aralia, fluffy ruffles fern, fuchsia, geranium, gerbera, gloxinia, heartleaf philodendron, hydrangea, kalanchoë, lantana, lily, marigold, pelargonium, petunia, purple passion plant, red-veined prayer plant, Sanders dracaena, snapdragon, table fern, variegated oval leaf peperomia, wax plant, weeping fig), foliage plants (areca palm, aucubifolium [cotton], coleus varieties, commutatum evergreen, coral ardisia, grape ivy, neanthe belle palm, sprengeri fern, zebra plant) without adverse effects.

Application of ENSTAR II should be made to poinsettia prior to bract formation. Foliar damage on some sensitive varieties can result; therefore, test on a small area before large-scale use. Some varieties of roses, such as yellow blooded roses, show delayed damage.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL (PLASTIC)

Triple rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Seller makes no warranty, express or implied, concerning the use of the product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

For information or in case of an emergency, call 1-800-248-7763 or visit our Web site: www.enstar2.com



Wellmark International
Shaumburg, Illinois U.S.A.

Enstar II A Wellmark International Brand.
ENSTAR® II Insect Growth Regulator and MAVERIK AQUAFLOW® are
registered trademarks of Wellmark International. ESS® is a registered
trademark of Electrostatic Spraying Systems. Puls-Fog® is a
registered trademark of Dramm International.